16.9" | 4.9" | 40.9" | 16.9" | 4.9"

29.00"

The Andersons®
Professional Turf™ Products

0.48% Barricade® on DG PRO®

Dispersing Granule Technology

GENERAL INFORMATION

This product is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses (excluding golf course putting greens), lawns, sod nurseries, landscape ornamentals, established perennial & wildflower plantings, non-crop areas on or surrounding managed rights-of-way for transportation systems and utilities (including roadways, roadsides, railways & equipment yards); facilities including substations, tankfarms, pumping stations, parking & storage areas, ungrazed fencerows.

This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Do not graze or feed livestock forage cut from areas treated with this product. Do not apply this product through any type of irrigation system unless otherwise instructed by this label. Do not apply aerially. Do not apply to golf course putting greens.

APPLICATION DIRECTIONS

Apply uniformly through suitable, calibrated application equipment.

ESTABLISHED TURF

This product is a preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the Maximum Annual Rates section of this label.

Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

CRABGRASS GERMINATION DATES



Approx. Date:
1 - After May 10
2 - After May 10
3 - After April 20
4 - After March 20
5 - Jan. 1 to March 20

Rates of Application

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

Maximum Annual Rates

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Application Rate of this product per Calendar Year by Turf Grass Species[1]			
	Lbs. AI/ A	Lbs. Product / A	Lbs. product/ 1,000 sq. ft.
Turf Species			
Bermudagrass[2], Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turf-type), Zoysia	1.5	312	7.1
Buffalograss, Kentucky Bluegrass, Perennial Ryegrass	1.0	208	4.7
Creeping Red Fescue	0.75	156	3.5
Creeping Bentgrass	0.65	135	3.1

[1]These are the maximum rates per calendar year by species limitations.
[2]May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 108 lb./acre or 2.4 lb./1,000 sq. ft. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Fontail, Goosegrass and Ressuegrass due to reduced product rates used in sprigging situations.

Use higher rates of this product to achieve higher levels weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates table.

Weeds Controlled

When used at the recommended rates this product will control:

Barnyardgrass	Goosegrass[1]	Pigweed
Bluegrass, Annual (Poa annual)	Henbit	Purslane, Common
Carpetweed	Itchgrass	Pusley, Florida
Chickweed, Common	Johnsongrass (from seed)	Rescuegrass[2]
Chickweed, Mouseear (from seed)	Junglerice	Shepherd's Purse[2]
Chickweed, Common (Large-flower)	Knotweed	Signalgrass, Broadleaf
Cinquefoil	Kochia	Speedwell, Persian
Crabgrass (Large, Smooth)	Lambsquarter, Common	Sprangletop
Crabgrass, Woolly	Lovegrass	Spurge, Prostrate
Foxtails, Annual	Oxalis	Witchgrass
	Panicum (Texas, Fall, Browntop)	Woodsorrel, Yellow (from seed)

[1]In many areas a single application of 0.1 to 7.1 lbs./1,000 sq.ft. (135-312 lbs./A) of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an application of 0.1 to 3.1 lbs./1,000 sq.ft. (135-208 lbs./A) followed after 60 to 90 days by a application at doses that would not exceed those given in the Maximum Annual Application Rates Table. Do not exceed the maximum rate for turf grass species listed in the Maximum Application Rates Table above.

[2]Applications for this weed should be made in late summer, fall, or winter prior to germination.

[3]Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

When to Apply After Overseeding Turf

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

When to Overseed After Application - All States*

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

The Andersons Professional Turf Products 0.48% Barricade on DG Pro			Interval Before Overseeding		
Lbs/1,000 sq. ft.	Lbs. /A	Ai/A	North	Transition	South
2.3	104	0.50	4 mo	4 mo	4 mo
3.1	135	0.66	5 mo	4 mo	4 mo
3.5	156	0.74	6 mo	5 mo	5 mo
3.8	166	0.80	---	6 mo	6 mo
4.7	208	1.00	---	7 mo	7 mo
5.4	237	1.14	---	---	9 mo
6.1	270	1.30	---	---	10 mo
7.1	312	1.50	---	---	12 mo

*Note: In AZ, CA, NV and TX, the overseeding interval can be shorter in established bermudagrass that has been overseeded with perennial ryegrass. See the next section "Poa annua Control in Established Bermudagrass Overseeded with Perennial Ryegrass (AZ, CA, NV and TX only)".

Poa annua Control in Established Bermudagrass Overseeded with Perennial Ryegrass (AZ, CA, NV and TX only)

Use on golf courses (excluding golf course putting greens) and lawns when overseeding with perennial ryegrass (minimum seed rate of 350 lbs./A).

How Much and When to Apply

Amount to Apply		When to Apply	Expected Control
Lbs./Acre	Lbs/1,000		
78 – 135	1.8 – 3.1	6 to 8 weeks before ryegrass overseeding	One application for 70% or greater control of *Poa annua*
		Second application 4-8 weeks after over-seeding date or when perennial ryegrass roots are in the second inch of soil	Second application should enhance the residual under normal weather conditions

Use Precautions

1. Some seed mortality and temporary reduction in root growth of new seedlings may occur.

2. To reduce the potential for seeding mortality, maintain a moist seedbed with light, frequent irrigation.

3. Make no more than 2 applications per year for this use, and do not exceed 4 lbs./1,000 sq. ft. (156 lbs./A) for these two applications for *Poa annua* control in ryegrass overseeded turf.

4. Do not make a second application if any injury to the ryegrass is observed after the first application.

5. Do not make a second application unless the product was first applied before overseeding.

Use Precautions

The following precautions apply to the use of this product in turf grasses and lawns:

Application of this product may thin emerged annual bluegrass and newly over seeded grasses. Do not apply to over seeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil. Do not apply to newly set sod until the following year. Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control. Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (*Poa annua*) are desirable species.

LANDSCAPE ORNAMENTALS

This product may be applied for residual preemergence weed control in ornamentals (including established perennial and wildflower plantings).

Use Rates

Apply this product at 3.1 to 7.1 lbs./1,000 sq. ft. (135 to 312 lbs./A) in fall and/or spring. Use higher rate of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 7.1 lbs./1,000 sq. ft. (312 lbs./acre).

Application Timing and Information

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves and mulch. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

Apply this product to dry foliage only. Overhead irrigation or rainfall soon after application will wash granules off plant foliage and activate this product in the soil.

Tolerant Ornamental Species

This product will not harm most trees, shrubs, vines, and flowers. The species below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface. This product may be applied over the top of these species.

When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

Fir species** (Balsam, Fraser, Noble, etc.)	Dianthus, Maiden Pinks 'Zing'
Abelia: Sherwood	Cheddar Pink
Japanese Maple	Hop Bush*
Norway Maple	Coneflower, Purple; Magnus
Yarrow: King Edward	Silverberry
Kiwi*	Wintercreeper
Lily-of-the-Nile (African Lily)	Japanese Spindle Tree (Evergreen Euonymus)
Five-Leaf or Chocolate Vine	Spreading Euonymus
Lady's Leek, Nodding Onion	Wintercreeper
Japanese Anemone	Border Forsythia
Aquilegia: Red and gold	Weeping Forsythia
Vine Hill Manzanilla	Greenstem Forsythia
Cape Weed	Gaillaria, Blanket Flower: 'Goblin'
Wormwood, Silver Mound, Castle	Gardenia, Cape-Jasmine; Radicans
Japanese Aucuba	Gentian*
Barberry	Cranesbill
Aster: Bonny Blue, Purple Dome	Gladiolus species**
Lady Fern; Fern Lady	Baby's Breath
Fibrous Begonia; Hardy Grandis	English Ivy
Wintergreen Barberry	Sunrose
Mentor Barberry	Daylily: Aztec Gold, Stella De Oro,
Japanese Barberry	Tender Love
Warty Barberry	Coral Bell; Bridget Bloom
Snowbank	Rose of Sharon**
Bougainvillea	Chinese Hibiscus
Butterfly Bush (Dwarf Blue); Royal Red	Mallow: Disco Belle White
Japanese Boxwood	Hosta, Plantain Lily (Fragrant)
Common Bottlebrush	Hosta, 'Searsucker'
Weeping Bottlebrush	Bigleaf Hydrangea
Stoplo Heather	Chinese Holly
Tussock Bellflower; (White Clips)	Japanese Holly; Helleri
Trumpet Creeper; Trumpet Flower; Madame Galen	American Holly
Hottentot Fig (Ice Plant)	Hosta
Feathery Cassia	Yaupon Holly; Schillings
Wild Lilac	Sword-Leaved Inc; Jodiesong
False Cypress	Siberian Iris; Cabernet
Cineyra	Iris species**
Citrus species*	Winter Jasmine
Coreopsis (Calliopsis); Early Sunrise	Chinese Juniper; Nick's Compact, Pyramid
Moonbean	Shore Juniper; Blue Pacific
Flowering Dogwood	Parsoni
American Dogwood	Creeping Juniper
Pampas Grass	Walnut*
Cranberry Cotoneaster	Shrimp Plant
Cotoneaster	Crape Myrtle
Bearberry Cotoneaster	Crape Myrtle: Tuscarora
Rockspray Cotoneaster	Weeping Lantana
Hawthorn	Lavender: Munstead
Lucifer	Edelweiss
Italian Cypress	Chinese Privet; Variegata
White Trailing Ice Plant	Amur Privet
Cooperi Pink	Japanese Privet
Larkspur: Blue Elf	Oleosy Privet (Wax-Leaf)
	Lily, Asiatic
	Big Blue Lillyturf; Turf, Silver Mound, Evergreen Giant
	Liriope, Variegated
	Liriope, Creeping
	Cardinal Flower, Indian Pink
	Japanese Honeysuckle
	Tatarian Honeysuckle
	Burgundy
	Loosestrife, Modern Pink
	Magnolia species**
	Ice Plant
	Crabapple*
	Yaku Jima, Silberfieder
	Heavenly Bamboo
	Narcissus species**
	Oleander
	Evening Primrose
	Olive*Mondo Grass
	Osmanthus (False Holly); Gulf Tide
	Trailing African Daisy
	Sourwood
	Tree Peony
	Fountain Grass (Dwarf)
	Avocado*
	Fraser's Photinia (Redtip)
	Dragonhead, False; Vivid
	Spruce species** (Colorado Blue, Dwarf Alberta)
	Lily-of-the-Valley Shrub
	Calabrian Pine
	Canary Island Pine
	Slash Pine
	Aleppo Pine
	Austrian Black Pine
	Longleaf Pine
	Monterey Pine
	Japanese Black Pine
	Eastern White Pine
	Scotch Pine
	Loblolly Pine
	Virginia Pine
	Pistachio*
	Queensland Pittosporum
	Japanese Pittosporum
	Japanese Yew
	English Laurel
	Almond, Apricot, Nectarine, Peach, Plum and Prune*
	Douglas Fir**
	Firethorn Scarlet
	Firethorn
	Bradford Pear spp.
	Oak, Shumard's Red
	Oak species
	Indian Hawthorn
	Yeddo Hawthorn
	Rhododendron (including Azalea)
	'Coral Bells'
	'Delaware Valley White'
	'Flame Creeper'
	'Formosa'
	'Girard Crimson'
	'George I. Tabor'
	'Hino-crimson'
	'PJM'
	'Roseum Elegans'
	Wakeiebisu
	White Gumbo
	Lady Bank's Rose
	Rosemary*
	Black-Eyed Susan; Goldstrum
	Leatherleaf Fern
	Santolina
	Saxifrage: Purple Dome
	Pincushion Flower
	Stonecrop; Lidakense, Dragon's Blood
	Spiraea: Anthony Waterer
	Australian Brushcherry
	Yew
	Germander
	Meadow Rue
	American Arborvitae
	Star Jasmine
	Canada Hemlock
	Tulip species
	Veronica Speedwell; Sunny Border
	Japanese Viburnum
	Sweet Viburnum
	Japanese Snowball
	Canary Island Viburnum
	Arrowood Viburnum
	Laurustinus
	Cranberry Bush
	Leatherleaf Viburnum
	Greater Periwinkle, Vinca
	Common Periwinkle, Vinca
	Grape*
	Old Fashioned Weigela
	Spanish Bayonet, Yucca
	Adam's Needle, Yucca

*Do not use on food producing trees, vines, or plants.
**Not for use on container grown plants.

VEGETATION MANAGEMENT

This product may be applied to soil surfaces for preemergence control of many grass and broadleaf weeds in non-cropland areas on or surrounding:

- Non-crop areas on or surrounding managed rights-of-way for transportation systems and utilities (including roadways, roadsides, railways & equipment yards)
- Facilities including substations, tankfarms, pumping stations, parking and storage areas and ungrazed fencerows

Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Apply this product at 3.1 to 7.1 lbs./1,000 sq. ft. (135 to 312 lbs./A) in fall and/or spring. Use higher rate of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 7.1 lbs./1,000 sq. ft. (312 lbs./A).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.
Storage: Store in original container away from fertilizer, feed, or food stuffs and separated from other pesticides.
Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty container into application equipment. Then dispose of empty bag or box in a sanitary landfill or at an incineration facility; or, if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE:
Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:
Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:
THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS. YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Product of U.S.A.

Visit our website: AndersonsLawnProducts.com

Manufactured by:
The Andersons Lawn Fertilizer Division, Inc.
P. O. Box 119
Maumee, OH 43537

©2008 The Andersons, Inc.
The Andersons logo and FOPro are registered trademarks of The Andersons, Inc.; Professional Turf is a trademark of The Andersons, Inc.
Barricade is a registered trademark of Syngenta Crop Portection

APTDG48BR5.1
APTDG48BR5.2B

FPO
UPC:
0 - 31865 - 31085 - 0

RECOMMENDED SPREADER SETTINGS

These suggested spreader settings are not intended to replace calibration. Calibrate your spreader before applying product.
50 lbs. treats 16,100 sq. ft. (0.37 acre) at the Low Rate (3.1 lbs./1,000 Sq. Ft.)
50 lbs. treats 14,200 sq. ft. (0.33 acre) at the Medium Rate (3.5 lbs./1,000 Sq. Ft.)
50 lbs. treats 10,600 sq. ft. (0.24 acre) at the High Rate (4.7 lbs./1,000 Sq. Ft.)
50 lbs. treats 7,000 sq. ft. (0.16 acre) at the Heavy Rate (7.1 lbs./1,000 Sq. Ft.)

		SPREADER SETTINGS			
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	LOW RATE 3.1 lbs./1,000 Sq. Ft.	MEDIUM RATE 3.5 lbs./1,000 Sq. Ft.	HIGH RATE 4.7 lbs./1,000 Sq. Ft.	HEAVY RATE 7.1 lbs./1,000 Sq. Ft.
Andersons AccuPro 2000, SR-2000 (cone 6)	3 mph	11 feet	K 1/2	L 1/2	M 1/2	P
Andersons SS-2 Drop	3 mph	overlap wheels	4 3/4	5	5 1/2	6 1/4
Scotts Rotaries: R-8, R-8A (cone 6)	3 mph	11 feet	J 1/2	K	L 3/4	O
LESCO 020093 (.25)	3 mph	5 feet	D 1/2	E	E 3/4	G 1/2
Lely Models WTR, WFT, HR, 1250 (PTO at 450 rpm) (Pattern IIIC)	4.5 mph	34 feet	5	5 1/4	6 1/4	7 1/4
Prizelawn CB85 B	3 mph	5 feet	3 1/2	3 3/4	4 1/4	4 3/4
Vicon (03 series)	4.5 mph	20 feet	17	18	22	28

The Andersons®
Professional Turf™ Products
0.48% Barricade® on DG PRO®
Product Code: APTDG48BR5.1

The Andersons®
Professional Turf™ Products
0.48% Barricade® on DG PRO®
Product Code: APTDG48BR5.1

The Andersons®
Professional Turf™ Products

0.48% Barricade® on DG PRO®

Pre-Emergent Weed Control for Lawns & Landscape Ornamentals

Dispersing Granule Technology

For preemergence control of grass and broadleaf weeds in established turfgrasses (excluding golf course putting greens), lawns and sod nurseries; landscape ornamentals; established perennial and wildflower plantings; non-crop areas on or surrounding managed rights-of-way for transportation systems and utilities (including roadways, roadsides, railways and equipment yards); facilities including substations, tankfarms, pumping stations, parking and storage areas, and ungrazed fencerows.

ACTIVE INGREDIENT:

Prodiamine	0.48%
OTHER INGREDIENTS:	99.52%
Total	100.00%

*OAS No. 29091-21-2)
EPA Reg. No. 9198-214
EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A
Underlined letter is first letter used in run code on bag

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes moderate eye injury irritation. Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE)
WPS USES:
Applicators and other handlers:
- Long sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash all PPE separately from other laundry.

NON-WPS USES:
Mixers and loaders who handles this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) – in general, only agricultural-plant uses are covered by the WPS – must wear:
- Chemical-resistant gloves

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash all PPE separately from other laundry.

Engineering Controls:
When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:
Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- After handling this product, immediately wash the outside of gloves before removing them, then remove gloves and all other PPE. Immediately wash thoroughly and change into clean clothing.

FIRST AID
If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

Environmental Hazards
This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations substantially above the level of water solubility, it may be toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

If material is spilled, contain and sweep up material of spill for disposal or reuse per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under or cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.

DIRECTIONS FOR USE
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS
Use this product only in accordance with its labeling and with the Workers Protection Standard 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:
- Coveralls
- Waterproof gloves
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS
The requirements in this box apply to use of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The Worker Protection Standard applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dusts have settled.

ACCEPTED
VIA NOTIFICATION
LABEL NOT REVIEWED
3/4/10
New York State Department of Environmental Conservation
Division of Solid & Hazardous Materials
Pesticide Product Registration

DOC ID 521642

Visit our website: AndersonsLawnProducts.com

Net Wt. 50 Lbs. (22.68 kg)

The Andersons®
Professional Turf™ Products
0.48% Barricade® on DG PRO®
Product Code: APTDG48BR5.1

FPO
UPC:
0 - 31865 - 31085 - 0

Product Code: APTDG48BR5.1

0.48% Barricade® on DG PRO®
The Andersons®
Professional Turf™ Products