

# LABEL SCAN CONTROL SHEET

---

Document ID 66035

Product ID 41140

EPA # 9198-184

Accepted  
Date 9/16/2003

ACCEPTED  
FOR REGISTRATION

SEP 16 2003

# K-O-G Weed Control

NEW YORK STATE DEPARTMENT  
OF ENVIRONMENTAL CONSERVATION  
DIVISION OF SOLID & HAZARDOUS MATERIALS  
Pesticide Product Registration

Record # 41140

•provides postemergence weed control

**Net Weight 35.50 lbs. (16:10 kg)**  
**Covers up to 44,000 Sq. Ft.**

**KEEP OUT OF REACH OF CHILDREN**  
**WARNING**  
**PRECAUTIONARY STATEMENTS**  
**Hazards to Humans and Domestic Animals**

**WARNING:** causes substantial but temporary eye injury. Harmful if absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Wear goggles or face shield. Harmful if swallowed. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Avoid inhalation.

**FIRST AID**

- If in eyes:**
  - Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing.
  - Call a poison control center or doctor for treatment advice.
- If on skin or clothing:**
  - Take off contaminated clothing.
  - Rinse skin immediately with plenty of water for 15-20 minutes.
  - Call a poison control center or doctor for treatment advice.
- If swallowed:**
  - Call a poison control center or doctor for treatment advice.
  - Have person sip a glass of water if able to swallow.
  - Do not induce vomiting unless told to by a poison control center or doctor.
  - Do not give anything by mouth to an unconscious person.
- If inhaled:**
  - Move person to fresh air.
  - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible.
  - Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

**Environmental Hazards:** Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

<b>ACTIVE INGREDIENT:</b>		
Dicamba (3,6-Dichloro-o-anisic acid)	0.70%	
<b>INERT INGREDIENTS:</b>	99.30%	
	Total	100.00%

EPA Reg. No. 9198-184  
EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A  
Underlined letter is first letter used in run code on bag.

Product of U.S.A.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Apply any time weeds are actively growing. Do not apply when ground is frozen.

**For bentgrass putting greens**

For bentgrass putting greens apply at LIGHT RATE to moist foliage for control of clover and chickweed. Repeat treatment in 3 weeks if needed. (Treats 44,000 sq. ft.)

**For use on other turfgrass areas**

For turfgrass areas other than greens, apply to moist or dry foliage at—

LOW RATE to control clover (treats 22,000 sq. ft.)

MEDIUM RATE to control knotweed, chickweed (common and mouse-ear), sheep sorrel and black medic (treats 11,000 sq. ft.)

HIGH RATE to control wild onion and garlic, curly dock, ground ivy, Canada thistle, cudweed, yarrow and horseweed. For control of yarrow and wild onion and garlic start treatments in either spring or fall and repeat at 6 month intervals for a total of 2 or 3 applications as needed (treats 5,500 sq. ft.)

**PRECAUTIONS**

Avoid application on rootzone areas of trees and shrubs.

Do not apply when ground is frozen.

Delay planting grass seed until after a heavy rain or watering and not sooner than two weeks after treatment.

On new plantings delay treatment until after the second mowing.

Not for use on St. Augustinegrass or Dichondra.

Avoid application to garden plants, flowers, and rootzone areas.

For use on fine fescue or bentgrass turf maintained at a height of 1/2 inch or above, use at the LOW RATE or LIGHT RATE setting.

Do not apply to slopes next to lakes or waterways or on slopes above beds used for trees or ornamental plantings.

Do not apply this product to turf under stress from drought, insects, disease, cold temperatures, high temperatures of above 85°F on cool season grasses, or poor fertility, as injury may result.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in a clean, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**IMPORTANT: READ BEFORE USE:**

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

**WARRANTY DISCLAIMER and LIMITATION OF LIABILITY:**

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

**LEGAL RIGHTS:**

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Manufactured by:  
The Andersons Lawn Fertilizer Division, Inc.  
P.O. Box 119  
Maumee, OH 43537

**SUGGESTED SPREADER SETTINGS**

Suggested spreader settings are not intended to replace calibration. Please calibrate your spreader prior to applying product.


- 35.5 lbs feeds 44,000 sq. ft. (1.01 acre) with 0.25 lbs. A/acre at the LIGHT RATE
- 35.5 lbs feeds 22,000 sq. ft. (0.50 acre) with 0.49 lbs. A/acre at the LOW RATE
- 35.5 lbs feeds 11,000 sq. ft. (0.25 acre) with 0.98 lbs. A/acre at the MEDIUM RATE
- 35.5 lbs feeds 5,500 sq. ft. (0.12 acre) with 1.96 lbs. A/acre at the HIGH RATE

	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS			
			LIGHT RATE	LOW RATE	MEDIUM RATE	HIGH RATE
The Andersons AccuPro/ SR-2000 (cone 9)	3 mph	10 feet	-	H 1/2	J 1/2	M 1/2
Scotts Rotaries R-8, R-8A (Cone 9)	3 mph	9 feet	-	G 1/2	I 1/2	L 1/2
Lely Models WTR, WFR, HR, 1250. (PTO at 450 rpm)	4.5 mph	26 feet	-	3 1/2 III-A	4 1/2 III-A	6 1/4 III-A
The Andersons SS-2 Drop	3 mph	-	2 1/2	3 1/4	4 1/4	5 1/2
Vicon (03 Series)	4.5 mph	16 feet	-	11	14	21

G3 (AGCBLKPB RN.1B)

**AGC8695**

*Logo that appears  
on blank (pocket) bag.*



**Andersons<sup>TM</sup>**  
**GOLF PRODUCTS**